

Abstract style for ISTCP-VII

Alice Waseda,¹ Bob Tokyo,² Charlie D. Author^{2,3}

¹*School of Advanced Science and Engineering, Waseda University, Japan*

²*Institute for Molecular Science, Japan*

³*Graduate School of Engineering, The University of Tokyo, Japan*

I. Introduction

The Seventh Congress of the International Society for Theoretical Chemical Physics (ISTCP-VII) will take place on September 2-8, 2011 at Waseda University (Fig. 1), Tokyo, Japan. A Congress is organized by ISTCP every three years. The venues of the past Congresses are listed in Table 1.

II. Preparation for abstract

The abstract should be no more than one page long when using a 12-point font on an A4 or letter-size paper [1] with margins of 35 mm from the top and 30 mm from the other sides. Please use Times New Roman (or Times / Roman) font through the abstract except for the title where 14-point Arial (or Helvetica / Sans Serif) font is used. Abstract title, author(s), and affiliation(s) should be centered. Please underline the presenting author. Affiliations should be given in italic letter. Although color figures are permitted in the electronic abstract, they are printed in black-and-white in the Abstract Book.

The abstract must be submitted in .pdf file format (.doc and .tex formats are prohibited). Please don't protect the PDF file because the presentation code will be filled electronically after the submission. The electronic submission website is accessible from the Abstract page of the Congress website [2].

III. Abstract submission deadlines

- Abstracts for Plenary, Invited, and Contributed Talks: March 31, 2011.
- Abstracts for Poster Presentation: June 30, 2011.

Table 1. ISTCP congress sites.

	Place	Year	Country
I	Girona	1993	Spain
II	New Orleans	1996	USA
III	Mexico City	1999	Mexico
IV	Paris	2002	France
V	New Orleans	2005	USA
VI	Vancouver	2008	Canada
VII ^a	Tokyo	2011	Japan

^a The first ISTCP congress held in Asia.



Figure 1. Okuma Auditorium, the symbol of Waseda University.

[1] A4: 210 mm × 297 mm, letter-size: 215.9 mm × 279.4 mm

[2] <http://www.chem.waseda.ac.jp/nakai/istcp7/>